

Applicant: Edward R. Burns and Vadiraja Murthy
Serial No.: 08/746,635
Filed: November 13, 1996
Page 2

Amendments to the Claims:

Please cancel claims 25 and 27-32 without disclaimer or prejudice to applicants' right to pursue the subject matter of these claims in a future continuation application. Please amend claims 20, 24 and 26 as set forth below.

1-19. (Canceled)

20. (Currently amended) A method for diagnosing erythrocyte hemolysis in a subject who is suspected of having ~~may have~~ erythrocyte hemolysis, the method comprising the steps of:

- (a) obtaining a serum sample from said subject; and
- (b) determining the level of erythrocyte adenylate kinase activity in said sample, the presence of at least about 20 U/L erythrocyte adenylate kinase activity in said sample being indicative of erythrocyte hemolysis in said subject.

21-23. (Canceled)

24. (Currently amended) ~~The method of claim 20, wherein determination of erythrocyte adenylate kinase activity comprises the steps of:~~

A method for diagnosing erythrocyte hemolysis in a subject who may have erythrocyte hemolysis, the method comprising the steps of:

- (a) obtaining a serum sample from the subject; and
- (b) determining the level of erythrocyte adenylate kinase activity in the sample by determining a value for total adenylate kinase activity in the serum sample;

299480.1

Applicant: Edward R. Burns and Vadiraja Murthy
Serial No.: 08/746,635
Filed: November 13, 1996
Page 3

determining a value for percentage erythrocyte adenylate kinase in the sample;
and

determining erythrocyte adenylate kinase activity in the sample by multiplying the value for percentage erythrocyte adenylate kinase by the value for total adenylate kinase activity,

the presence of at least about 20 U/L erythrocyte adenylate kinase activity in the sample being indicative of erythrocyte hemolysis in the subject.

25. (Canceled)

26. (Currently amended) ~~The method of claim 25, wherein the antibody is specific for erythrocyte adenylate kinase~~

A method for diagnosing erythrocyte hemolysis in a subject who may have erythrocyte hemolysis, the method comprising the steps of:

(a) obtaining a serum sample from the subject; and

(b) determining the level of erythrocyte adenylate kinase activity in the sample using an antibody that is specific for and binds to erythrocyte adenylate kinase,

the presence of at least about 20 U/L erythrocyte adenylate kinase activity in the sample being indicative of erythrocyte hemolysis in the subject.

27-32. (Canceled)

33. (Previously presented) The method of claim 26, where the antibody is detected using a radioimmunoassay.

299480.1

Applicant: Edward R. Burns and Vadiraja Murthy
Serial No.: 08/746,635
Filed: November 13, 1996
Page 4

34. (Previously presented) The method of claim 26, where the antibody is detected using an enzyme-linked immunosorbent assay (ELISA).

35. (Previously presented) The method of claim 26, where the antibody is detected using immunoprecipitation.

36. (Previously presented) The method of claim 26, wherein the antibody is labeled with a detectable marker.

37. (Previously presented) The method of claim 26, wherein the antibody is labeled with a radioisotope.

38. (Previously presented) The method of claim 26, wherein the antibody is labeled with radioactive iodine.